arXiv:1208.4917v1 [cond-mat.supr-con] 24 Aug 2012

Single nonmagnetic impurity resonance in FeSe-based 122-type superconductors as a probe for pairing symmetry

Qian-En Wang, Zi-Jian Yao, and Fu-Chun Zhang
Department of Physics and Center of Theoretical and Computational Physics,
The University of Hong Kong, Hong Kong, China

(Dated: August 27, 2012)

The effect of a single nonmagnetic impurity in $A_yFe_{2-x}Se_2$ (A=K, Rb, or Cs) superconductors has been studied based on a three-orbital model. The local density of states on and near the impurity site has been calculated by solving the Bogoliubov-de Gennes equations self-consistently. Both repulsive and attractive impurity scattering potential are considered in our calculations. The impurity-induced in-gap bound states are found only for attractive scattering potential in the state of $d_{x^2-y^2}$ wave pairing, and it turns out that they are very sensitive to the magnitude of the scattering potential. The emergence of the impurity-induced bound states in the vicinity of the Fermi level demonstrates a strong violation of the electron-hole symmetry which is originated from the nodeless $d_{x^2-y^2}$ wave pairing state. The results obtained in our calculation, which simulate the doping of Co and Ni in FeSe-based 122-type superconductors, as an approach to examine the pairing symmetry of this novel superconducting material, can be a proposal of STM observation.

PACS numbers: 74.70.Xa, 74.25.Jb, 74.20.Pq

I. INTRODUCTION

The FeSe-based 122-type $A_yFe_{2-x}Se_2$ (A=K, Rb, or Cs) high-T_c superconductors have been focused intensively due to its distinctive high transition temperatures(∼30K)[1–4] and novel Fermi surface topology. Angle-resolved photoemission spectroscopy (ARPES)[5–9] and density functional theory (DFT) calculations[10–13] show that there is no hole pocket around the Γ point of the Brillouin zone (BZ), and large electron pockets are observed around the M point which is located at the corners of the folded BZ[5]. Effort has been made to investigate the pairing symmetry of the Fe-based superconductors. The popular $s_\pm$ pairing symmetry in iron pnictide superconductors in which the hole pockets are at the Γ point, has been predicted by spin-fluctuation theory[14,15]. However, for FeSe-based 122-type superconductors, due to the absence of the hole pockets, nodeless superconducting (SC) gap can be originated from various pairing symmetries. Spin-fluctuation theory predicted $d_{x^2-y^2}$ wave pairing state[16–19], while s wave pairing state was argued to be more favorable against strong disorder[20]. Moreover, a co-existence candidate of anisotropic s and $d_{x^2-y^2}$ wave pairing symmetries has also been predicted[21]. These results reveal the competitive nature of the pairing state in these novel superconductors.

The response of a SC system to the impurities can be a probe for the SC pairing symmetry[22]. The local electronic states can be influenced by impurities evidently. In a multi-band SC system, the SC order parameter may or may not be sensitive to the existence of impurity[23]. Doping of Co and Zn in iron pnictide superconductors has been realized by several experimental groups[24–28]. Theoretically, the effects of impurity in iron pnictide superconductors has been investigated intensively[29–35]. The calculated results of T-matrix method based on a four-band model shows the in-gap impurity resonance states[33] for $s_\pm$ wave pairing symmetry, and results obtained by solving the Bogoliubov-de Gennes (BdG) equations in the framework of a detailed five-orbital model also demonstrates that the $s_\pm$ wave pairing symmetry is characterized by a bound state in the vicinity of the Fermi level in the presence of a single nonmagnetic impurity[34]. However, for FeSe-based 122-type superconductors, the impurity resonance bound state, which is located away from the Fermi level, were obtained only in $d_{x^2-y^2}$ wave pairing state[36].

In this work, we study the effect of a single nonmagnetic impurity in FeSe-based 122-type superconductors based on a three-orbital model and by using BdG[40] equations. The local density of states(LDOS) and the real-space SC order parameters are calculated. It has been reported that the effective on-site impurity potentials can be repulsive or attractive which is related to the energy difference between the impurity 3d levels and Fe 3d level for various impurity atoms. Realistic estimation of the impurity potential in iron pnictide superconductors predicted that the potential difference of Zn, Co, and Ni are negative, whereas Mn are positive[37]. Therefore both repulsive(positive) and attractive(negative) scattering potentials have been considered in our calculation which may simulate different dopants, such as Zn, Co, Ni, and Mn etc.. Responses of isotropic s, anisotropic s, and $d_{x^2-y^2}$ wave pairing states to impurity are examined, respectively. We focus on the difference of the resonance states induced by impurity among different pairing symmetries and the properties of the impurity-induced bound states.

This paper is organized as follows. Section II presents the Hamiltonian of the three-orbital model and the self-consistent BdG approach. In section III, we discuss and compare the results of the properties of the impurity states for different pairing symmetries. Finally, a summary is given in section IV.

II. MODEL AND METHODOLOGY

The Hamiltonian of the impure SC system is of the following form

$$H = H_0 + H_{pair} + H_{imp} \tag{1}$$

and the hopping, pairing, and impurity terms read

$$\begin{aligned} H_0 &= \sum_{i,j,\alpha,\beta,\sigma} t_{ij}^{\alpha\beta} a_{i\alpha\sigma}^{\dagger} a_{j\beta\sigma} - \mu \sum_{i,\alpha,\sigma} a_{i\alpha\sigma}^{\dagger} a_{i\alpha\sigma} \\ H_{pair} &= \sum_{i,j,\alpha,\beta} (\Delta_{ij}^{\alpha\beta} a_{i\alpha\uparrow}^{\dagger} a_{j\beta\downarrow}^{\dagger} + h.c.) \\ H_{imp} &= U_s \sum_{\alpha,\sigma} a_{l\alpha\sigma}^{\dagger} a_{l\alpha\sigma} \end{aligned} \tag{2}$$

where $a_{i\alpha\sigma}^{\dagger}(a_{i\alpha\sigma})$ denotes the creation(annihilation) operator of electrons with spin $\sigma = \uparrow, \downarrow$ and orbital α at site i. $t_{ij}^{\alpha\beta}$ are the hopping integrals and μ is the chemical potential. A single nonmagnetic impurity is located at site l. Results of DFT calculation demonstrates that the electronic structure in the vicinity of the Fermi level is dominated by d_{xz}, d_{yz}, and d_{xy} orbitals[13], therefore it is feasible to describe the band structure by introducing an effective three-orbital model[38]. The hopping parameters which are given in irreducible hopping subsets[18] are shown in the Appendix. Fig. 1 shows the heavily electron-doped Fermi surface obtained with a chemical potential μ=0.312 eV corresponding to a filling factor n=4.23. The SC gap function stemming from the mean-field decoupling of the pair scattering process $V_{ij}^{\alpha\beta} a_{i\alpha\uparrow}^{\dagger} a_{j\beta\downarrow}^{\dagger} a_{j\beta\downarrow} a_{i\alpha\uparrow}$ is expressed as

$$\Delta_{ij}^{\alpha\beta} = V_{ij}^{\alpha\beta} \langle a_{j\beta\downarrow} a_{i\alpha\uparrow} \rangle \tag{3}$$

Single static impurity in a uniform superconductor which manifests itself as a point-like imperfection can be regarded as a localized orbital-independent on-site scattering potential $U_s \delta_{\hat{R}_l R_0}$ with the impurity at R_0 in orbital representation, provided that the Coulomb interaction is screened at length scales comparable to the lattice spacing[22]. Both positive and negative scattering potential U_s are considered in our calculation, and by checking the results obtained from several different values, it has been found that $U_s = \pm 6.0$ eV can be regarded as the unitary limit.

The Hamiltonian of the impure system which is originally written in orbital representation is of its diagonal form in qusiparticle representation by conducting the Bogoliubov-Valatin transformation[40,41]

$$\begin{pmatrix} \gamma_{n\uparrow} \\ \gamma_{n\downarrow}^{\dagger} \end{pmatrix} = \sum_{i,\alpha} \begin{pmatrix} u_{i\alpha}^{n*} & v_{i\alpha}^{n*} \\ -v_{i\alpha} & u_{i\alpha}^{n} \end{pmatrix} \begin{pmatrix} a_{i\alpha\uparrow} \\ a_{i\alpha\downarrow}^{\dagger} \end{pmatrix} \tag{4}$$



FIG. 1: (color online) The Fermi surface in the extended BZ. The nodal lines of the gap function for anisotropic s and d$_{x^2-y^2}$ wave pairing symmetry are plotted with green and red lines, respectively.

where the quasiparticle creation operator $\gamma_{n\sigma}^{\dagger}$ is the ladder operator of the eigen-spectrum of the Hamiltonian which satisfies $[\gamma_{n\sigma}^{\dagger}, H]_- = -\epsilon_n \gamma_{n\sigma}^{\dagger}$.

The diagonal condition of the Hamiltonian is the BdG equation

$$\sum_{j,\beta} \begin{pmatrix} h_{ij}^{\alpha\beta} & \Delta_{ij}^{\alpha\beta} \\ \Delta_{ij}^{\alpha\beta *} & -h_{ij}^{\alpha\beta *} \end{pmatrix} \begin{pmatrix} u_{j\beta}^{n} \\ v_{j\beta}^{n} \end{pmatrix} = \epsilon_n \begin{pmatrix} u_{i\alpha}^{n} \\ v_{i\alpha}^{n} \end{pmatrix} \tag{5}$$

where

$$h_{ij}^{\alpha\beta} = t_{ij}^{\alpha\beta} - \mu \delta_{ij} \delta_{\alpha\beta} + U_s \delta_{il} \delta_{jl} \delta_{\alpha\beta}$$

and the gap equation is represented as

$$\Delta_{ij}^{\alpha\beta} = -\frac{V_{ij}^{\alpha\beta}}{2} \sum_n (u_{i\alpha}^{n} v_{j\beta}^{n*} + u_{j\beta}^{n} v_{i\alpha}^{n*}) \tanh(\frac{\epsilon_n}{2k_B T}) \tag{6}$$

The gap function of BCS theory[42] is an eigenvalue equation of pairing potential in k-space[39], the symmetric property of which is characterized by the irreducible representation of the symmetry group of the Hamiltonian. Therefore, the symmetry of the gap function is determined by the form of the pairing potential. The absence of the hole pockets of the FeSe-based 122-type Fermi surface around Γ point eliminates the possibility of nodes. Calculation pertaining to the magnetic exchange couplings reveals that the leading pairing is originated from the intra-orbital pairing, whereas the inter-orbital components are found to be significantly small[38]. Consequently, the s and $d_{x^2-y^2}$ wave pairing symmetries are considered for intra-orbital pairing only in order to investigate their responses to a single nonmagnetic impurity scattering. For isotropic s wave pairing,

$$V_{ij}^{\alpha\alpha} = -g_0 \delta_{ij} \tag{7}$$

for anisotropic s wave pairing,

$$\begin{aligned} V_{ij}^{\alpha\alpha} &= -\frac{g_1}{4}(\delta_{i+\hat{x}+\hat{y},j} + \delta_{i-\hat{x}+\hat{y},j} \\ &+ \delta_{i-\hat{x}-\hat{y},j} + \delta_{i+\hat{x}-\hat{y},j}) \end{aligned} \tag{8}$$

and for $d_{x^2-y^2}$ wave pairing,

$$\begin{aligned} V_{ij}^{\alpha\alpha} &= -\frac{g_2}{2}(\delta_{i+\hat{x},j} + \delta_{i+\hat{y},j} \\ &+ \delta_{i-\hat{x},j} + \delta_{i-\hat{y},j}) \end{aligned} \quad (9)$$

where $g_{0,1,2}$ are pairing amplitudes for each pairing symmetry and $\hat{x}(\hat{y})$ denote the unit vectors along $x(y)$ directions of the lattice coordinate. The pairing patterns in real space result in specific pairing symmetries of the gap function in k-space as for isotropic s wave channel $\Delta^{\alpha\alpha}(k) \sim \Delta_0$, for anisotropic s wave channel $\Delta^{\alpha\alpha}(k) \sim \cos(k_x)\cos(k_y)$, and for $d_{x^2-y^2}$ wave channel $\Delta^{\alpha\alpha}(k) \sim \cos(k_x) - \cos(k_y)$ in the extended BZ, respectively. The pairing amplitudes are taken to be $g_0 = 0.3$, $g_1 = 1.0$, and $g_0 = 0.5$ which guarantees that the SC coherence length $\xi \leq 4a$, where a is the lattice constant.

The energy spectrum of the quasiparticle i.e., the LDOS at site i is calculated via

$$\rho_i(\epsilon) = \frac{1}{N_{orb}} \sum_{n,\alpha} [|u_{i\alpha}^n|^2 \delta(\epsilon - \epsilon_n) + |v_{i\alpha}^n|^2 \delta(\epsilon + \epsilon_n)] \quad (10)$$

where the prefactor comes from taking the average of three orbitals. The Lorentzian smearing method is used to visualize the LDOS with a broadening width $\sigma = 0.001$. All the self-consistent calculations are performed on a 20×20 lattice with a periodic boundary condition at temperature T= 0.1K, and the single impurity is located at the center of the lattice.

III. RESULTS AND DISCUSSION

The calculated LDOS for isotropic s wave pairing state are shown in Fig. 2 for $U_s = 1.5$ eV and $U_s = -2.0$ eV, respectively. Obviously, the existence of impurity does not change the global properties of the quasiparticle energy spectrum. Similar results are also obtained in Ref.[29,34] which demonstrates that there is no in-gap bound state for such a pairing symmetry, whereas the order parameter on the impurity site are suppressed evidently.

Results of anisotropic s pairing symmetry are shown in Fig. 3 for repulsive scattering potentials and in Fig. 4 for attractive cases, respectively. The SC coherence peak is located at ±0.044 eV, and the typical resonating states induced by the impurity appear for both cases at ±0.028 eV for $U_s = 0.5$ eV and ±0.025 eV for $U_s = -1.0$ eV, respectively. The resonance peaks move towards the gap-edge as the amplitude of the scattering potential increases for repulsive interaction, while for attractive cases, the resonating peaks maintain at the gap-edge for small value of scattering potential $U_s = -0.5$ eV, as shown in Fig. 4(a). Additionally, it has been noticed that the resonating states for positive interaction $U_s = 0.5$ eV is very robust in comparison with other values of U_s. At unitary limit, the characteristics of the spectrum for both repulsive and attractive interaction tends to be identical since the electrons cannot hop to the impurity site, regardless of the detailed feature of the scattering interaction. We note that the resonating states within the SC gap, as compared with the clean cases, appear always in pairs which are located symmetrically with respect to the Fermi level. The magnitudes of the LDOS of the resonating states at $E = \pm 0.028(\pm 0.025)$ eV for $U_s = 0.5(-1.0)$ eV, for instance, may not be the same, but they are of the same order. Therefore, we may interpret them as the SC pairing states influenced by the impurity.

The response of $d_{x^2-y^2}$ wave pairing state to impurity scattering in cuprate superconductors has been investigated intensively, and the study of the impurity effects in superconductors has been considered as an effective method to reveal the physics of superconductivity[22]. Fig. 5 and 6 show the LDOS in the state of $d_{x^2-y^2}$ wave



(a) (b)

FIG. 2: (color online) The LDOS for isotropic s wave pairing state for $U_s = 1.5$ eV (a) and $U_s = -2.0$ eV (b). LDOS for impurity, NN, and NNN sites are plotted in red, green, and blue lines, respectively. The spectrum of the clean system is also shown in black line for comparison.



(a) (b)

(c) (d)

FIG. 3: (color online) The LDOS for anisotropic s wave pairing state for repulsive scattering potential $U_s = 0.5$ eV (a), $U_s = 0.8$ eV (b), $U_s = 2.0$ eV (c), and $U_s = 6.0$ eV (d).



(a) (b) (c) (d)

FIG. 4: (color online) The LDOS for anisotropic s wave pairing state for attractive scattering potential $U_s = -0.5$ eV (a), $U_s = -1.0$ eV (b), $U_s = -2.5$ eV (c), and $U_s = -6.0$ eV (d).



(a) (b)

FIG. 5: (color online) The LDOS for $d_{x^2-y^2}$ wave pairing symmetry for repulsive scattering potential $U_s = 0.5$ eV (a) and $U_s = 2.0$ eV (b).



(a) (b) (c) (d) (e) (f)

FIG. 6: (color online) The LDOS for d_{x2-y2} wave pairing state for attractive scattering potential $U_s = -0.5$ eV (a), $U_s = -0.8$ eV (b), $U_s = -1.0$ eV (c), $U_s = -1.5$ eV (d), $U_s = -2.0$ eV (e), and $U_s = -6.0$ eV (f).



(a) (b)

FIG. 7: (color online) Real-space distribution of the order parameter $\Delta(i)$ of anisotropic s (a) and $d_{x^2-y^2}$ (b) wave pairing states at unitary limit of the scattering interaction.

pairing for positive and negative scattering potential, respectively. It has been found that there is no in-gap bound state for repulsive scattering potential cases. However, for attractive scattering interaction, the impurity-induced in-gap bound states exhibit a high electron-hole asymmetry. The locations of the impurity states move towards the Fermi level from the positive energy as the amplitude of the scattering potential increases from 0.5 to 1.0 (eV), and shifts to the negative energy then to the gap-edge as the intensity of the potential approaches to the unitary limit. Obviously, these are impurity-induced bound states stemming from the sign-change of the $d_{x^2-y^2}$ wave SC gap. In our calculations, no evident in-gap bound states are observed when the scattering potentials are small enough as $|U_s| \leq 0.4$ eV, which is physically justified that the impurity-induced bound states would become small and vanish as the amplitude of scattering interaction decreases. Decomposed LDOS analysis demonstrates that the contribution of d_{xy} orbital is larger than that of d_{xz} and d_{yz} orbitals which indicates the significance of the multi-orbital feature of the Fe-based superconductors. The distribution of the order parameter in real-space at unitary limit of the impurity scattering, as plotted in Fig. 7, shows a color mapping of the cross shaped[22] suppression of the order parameters which occurs prominently along the NNN directions for anisotropic s wave and NN directions for $d_{x^2-y^2}$ wave pairing state, respectively.

Different from the SC pairing states which are influenced by the impurity in the state of anisotropic s wave pairing, the single impurity-induced in-gap bound state, for the $d_{x^2-y^2}$ wave pairing symmetry, is located near the Fermi energy for a moderate scattering potential $U_s = -1.0$ eV. This is reminiscent of the cuprates where the emergence of the impurity-induced bound states, as a result of the nodal $d_{x^2-y^2}$ wave pairing state, is characterized by its low-lying feature. It has been noticed that the amplitude of the density of the impurity-induced bound states is determined by the Bogoliubov quasiparticle weights $|u^n_{i\alpha}|^2$ and $|v^n_{i\alpha}|^2$, and the asymmetry of LDOS of the bound states in Fig. 6 demonstrates a strong violation of the electron-hole symmetry. Therefore, for FeSe-based 122-type superconductors, the nodeless $d_{x^2-y^2}$ wave pairing state leads to a peculiar bound states which are characterized by a strong electron-hole asymmetry. Particularly, only one single robust bound state on impurity site has been found in the gap for the attractive scattering interaction. This is consistent with the results obtained by Zhu *et al.*[36] for repulsive scattering potential. In our calculation, however, it has been found that there is no impurity-induced bound state for repulsive scattering interaction in the state of $d_{x^2-y^2}$ wave pairing, which can be interpreted as the effect of the strong violation of the electron-hole symmetry due to the peculiar Fermi surface of the FeSe-based 122-type superconductors where only electron pockets exist.

Additionally, the subtle mechanism of the substitution of Co and Ni for Fe remains a controversial question in Fe-based superconductors. The effective impurity potential induced by various dopants which can be ascribed to the difference of the on-site ionic potential, together with the modification of the Fermi surface caused by the rigid shift of the band structure, has been investigated from both theoretical and experimental perspective[37,43,44]. The isovalence of Co and Fe in Ca(Fe$_{0.944}$Co$_{0.056}$)$_2$As$_2$ compound has been pointed out and a binding energy shift (0.25 eV) has also been observed, which reveal the complexity of the impurity nature of Co[44]. Inspired by this finding, one may anticipate that the doping of Ni in iron pnictide or chalcogenide superconductors may be another promising alternative since the effective impurity potential, as reported by Nakamura *et al.*[37], is higher than that of Co. The impurity-induced bound states observed in our calculations in the state of $d_{x^2-y^2}$ wave pairing when the scattering potential is within the range of $U_s = -0.5 \sim -6.0$ eV may simulate the cases of Ni and Co doping in FeSe-based 122-type superconductors.

IV. SUMMARY

In summary, using a three-orbital model, we present a comprehensive investigation of the effects of a single nonmagnetic impurity in A$_x$Fe$_{2-y}$Se$_2$ (A=K, Rb, or Cs) superconductors by means of BdG theory. The in-gap bound states has been found for $d_{x^2-y^2}$ wave pairing state. For anisotropic s wave pairing state, the observed resonating states are typically SC pairing states influenced by impurity. The locations of the bound states for $d_{x^2-y^2}$ pairing symmetry, moving towards the Fermi energy from the gap-edge as the amplitude of the scattering potential increases from 0.5 to 1.0 (eV), appear uniquely for attractive interaction. Due to the absence of the hole pocket, strong violation of the electron-hole symmetry occurs, which results in that no impurity-induced bound state emerges for repulsive scattering interaction. Therefore, distinguishable responses to a single nonmagnetic impurity, in terms of LDOS on and near the impurity site, would be an approach to examine the pairing symmetries of FeSe-based 122-type superconductors.

Appendix: irreducible hopping matrices of the three-band model

The band structure of the three-orbital model is obtained by diagonalizing the Hamiltonian in orbital representation. The D_{4h} point group symmetry guarantees that the Bloch Hamiltonian can be written in a compact rotational form as the following

$$H = \sum_k \psi^\dagger(k) M(k) \psi(k) \tag{A.1}$$

where $\psi(k) = [a^\dagger_{xz}(k), a^\dagger_{yz}(k), a^\dagger_{xy}(k)]$ consists of the orbital-component operators and

$$\begin{aligned} M(k) &= K_0 \\ &+ K_1 e^{ik_x} + R K_1 R^{-1} e^{ik_y} \\ &+ R^2 K_1 R^{-2} e^{-ik_x} + R^3 K_1 R^{-3} e^{-ik_y} \\ &+ K_2 e^{i(k_x+k_y)} + R K_2 R^{-1} e^{i(-k_x+k_y)} \\ &+ R^2 K_2 R^{-2} e^{i(-k_x-k_y)} + R^3 K_2 R^{-3} e^{i(k_x-k_y)} \end{aligned} \tag{A.2}$$

where the point group operation is carried out by rotation

$$R = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \tag{A.3}$$

The irreducible hopping subsets (in unit: eV) corresponding to on-site atomic energies, hopping along $\hat{x}$, and $\hat{x}+\hat{y}$ directions are

$$\begin{aligned} K_0 &= \begin{pmatrix} 0.00 & 0.00 & 0.00 \\ 0.00 & 0.00 & 0.00 \\ 0.00 & 0.00 & 0.40 \end{pmatrix} \\ K_1 &= \begin{pmatrix} 0.05 & 0.00 & -0.20 \\ 0.00 & 0.01 & 0.00 \\ 0.20 & 0.00 & 0.20 \end{pmatrix} \\ K_2 &= \begin{pmatrix} 0.02 & 0.01 & 0.10 \\ 0.01 & 0.02 & 0.10 \\ -0.01 & -0.10 & 0.20 \end{pmatrix} \end{aligned} \tag{A.4}$$

For simplicity, the spin index has been dropped.

---

[1] Jiangang Guo, Shifeng Jin, Gang Wang, Shunchong Wang, Kaixing Zhu, Tingting Zhou, Meng He, and Xiaolong Chen, Phys. Rev. B **82**, 180520 (2010).

[2] Ming-Hu Fang, Hang-Dong Wang, Chi-Heng Dong, Zu-Juan Li, Chun-Mu Feng, Jian Chen, and H. Q. Yuan, Europhys. Lett. **94**, 27009 (2011).

[3] R. H. Liu, X. G. Luo, M. Zhang, A. F. Wang, J. J. Ying, X. F. Wang, Y. J. Yan, Z. J. Xiang, P. Cheng, G. J. Ye, Z. Y. Li, and X. H. Chen, Europhys. Lett. **94**, 27008 (2011).

[4] A Krzton-Maziopa, Z Shermadini, E Pomjakushina, V Pomjakushin, M Bendele, A Amato, R Khasanov, H Luetkens, and K Conder, J. Phys. Condens. Matter **23**, 052203 (2011).

[5] Y. Zhang, L. X. Yang, M. Xu, Z. R. Ye, F. Chen, C. He, H. C. Xu, J. Jiang, B. P. Xie, J. J. Ying, X. F. Wang, X. H. Chen, J. P. Hu, M. Matsunami, S. Kimura, and D. L. Feng, Nature Mater. **10**, 273 (2011).

[6] Daixiang Mou, Shanyu Liu, Xiaowen Jia, Junfeng He, Yingying Peng, Lin Zhao, Li Yu, Guodong Liu, Shaolong He, Xiaoli Dong, Jun Zhang, Hangdong Wang, Chiheng Dong, Minghu Fang, Xiaoyang Wang, Qinjun Peng, Zhimin Wang, Shenjin Zhang, Feng Yang, Zuyan Xu, Chuangtian Chen, and X. J. Zhou, Phys. Rev. Lett. **106**, 107001 (2011).

[7] X.-P. Wang, T. Qian, P. Richard, P. Zhang, J. Dong, H.-D. Wang, C.-H. Dong, M.-H. Fang, and H. Ding, Europhys. Lett. **93** 57001 (2011).

[8] T. Qian, X.-P.Wang, W.-C. Jin, P. Zhang, P. Richard, G. Xu, X. Dai, Z. Fang, J.-G. Guo, X.-L. Chen, and H. Ding, Phys. Rev. Lett. **106**, 187001 (2011).

[9] Lin Zhao, Daixiang Mou, Shanyu Liu, Xiaowen Jia, Junfeng He, Yingying Peng, Li Yu, Xu Liu, Guodong Liu, Shaolong He, Xiaoli Dong, Jun Zhang, J. B. He, D. M. Wang, G. F. Chen, J. G. Guo, X. L. Chen, Xiaoyang Wang, Qinjun Peng, Zhimin Wang, Shenjin Zhang, Feng Yang, Zuyan Xu, Chuangtian Chen, and X. J. Zhou, Phys. Rev. B **83**, 140508(R) (2011).

[10] Chao Cao and Jianhui Dai, Chin. Phys. Lett. **28**, 057402 (2011).

[11] Lijun Zhang and D. J. Singh, Phys. Rev. B **79**, 094528 (2009).

[12] Xun-Wang Yan, Miao Gao, Zhong-Yi Lu, and Tao Xiang, Phys. Rev. B **84**, 054502 (2011).

[13] I. R. Shein and A. L. Ivanovskii, Phys. Lett. A **375**, 1028-1031 (2011).

[14] Kazuhiko Kuroki, Seiichiro Onari, Ryotaro Arita, Hidetomo Usui, Yukio Tanaka, Hiroshi Kontani, and Hideo Aoki, Phys. Rev. Lett. **101** 087004 (2008).

[15] I. I. Mazin, D. J. Singh, M. D. Johannes, and M. H. Du, Phys. Rev. Lett. **101** 057003 (2008).

[16] Tanmoy Das and A. V. Balatsky, Phys. Rev. B **84** 014521 (2011).

[17] T. A. Maier, S. Graser, P. J. Hirschfeld, and D. J. Scalapino, Phys. Rev. B **83** 100515(R) (2011).

[18] Fa Wang, Fan Yang, Miao Gao, Zhong-Yi Lu, Tao Xiang, and Dung-Hai Lee, Europhys. Lett. **93** 57003 (2011).

[19] Tetsuro Saito, Seiichiro Onari, and Hiroshi Kontani, Phys. Rev. B **83** 140512(R) (2011)

[20] Yi Zhou, Dong-Hui Xu, Fu-Chun Zhang, and Wei-Qiang Chen, Europhys. Lett. **95** 17003 (2011).

[21] Rong Yu, Pallab Goswami, Qimiao Si, Predrag Nikolic, and Jian-Xin Zhu, arXiv: 1103.3259

[22] A. V. Balatsky, I.Vekhter, and Jian-Xin Zhu, Rev. Mod. Phys. **78**, 373 (2006).

[23] J. M. Byers, M. E. Flatté, and D. J. Scalapino, Phys. Rev. Lett. **71**, 3363 (1993).

[24] Ayaka KAWABATA, Sang Chul LEE, Taketo MOYOSHI, Yoshiaki KOBAYASHI, and Masatoshi SATO, J. Phys. Soc. Jpn. **77**, 103704 (2008).

[25] Ayaka KAWABATA, Sang Chul LEE, Taketo MOYOSHI, Yoshiaki KOBAYASHI, and Masatoshi SATO, J. Phys. Soc. Jpn. **77** Suppl. C, pp147 (2008).

[26] Athena S. Sefat, Rongying Jin, Michael A. McGuire, Brian C. Sales, David J. Singh, and David Mandrus, Phys. Rev. Lett. **101** 117004 (2008).

[27] A. Leithe-Jasper, W. Schnelle, C. Geibel, and H. Rosner, Phys. Rev. Lett. **101** 207004 (2008)

[28] Yuke Li, Xiao Lin, Qian Tao, Cao Wang, Tong Zhou, Linjun Li, Qingbo Wang, Mi He, Guanghan Cao, and Zhu'an Xu, New J. Phys. **11**, 053008 (2009)

[29] Tao Zhou, Xiang Hu, Jian-Xin Zhu, and C. S. Ting, arXiv: 0904.4273 (2009).

[30] Masashige MATSUMOTO, Mikito KOGA, and Hiroaki KUSUNOSE, J. Phys. Soc. Jpn. **78**, 084718 (2009).

[31] Tao Zhou, Huaixiang Huang, Yi Gao, Jian-Xin Zhu, and C. S. Ting, Phys. Rev. B **83**, 214502 (2011).

[32] Yuko Senga, and Hiroshi Kontani, New J. Phys. **11**, 035005 (2009).

[33] Degang Zhang, Phys. Rev. Lett. **103**, 186402 (2009).

[34] Toshikaze KARIYADO and Masao OGATA, J. Phys. Soc.

Jpn. **79**, 083704 (2010).

[35] Wei-Feng Tsai, Yan-Yang Zhang, Chen Fang, and Jiangping Hu, Phys. Rev. B **80**, 064513 (2009).

[36] Jian-Xin Zhu, Rong Yu, A. V. Balatsky, and Qimiao Si, Phys. Rev. Lett. **107**, 167002 (2011).

[37] Kazuma Nakamura, Ryotaro Arita, and Hiroaki Ikeda, Phys. Rev. B **83**, 144512 (2011)

[38] Chen Fang, Yang-Le Wu, Ronny Thomale, B. Andrei Bernevig, and Jiangping Hu, Phys. Rev. X **1**, 011009 (2011).

[39] M. Sigrist and K. Ueda, Rev. Mod. Phys. **63**, 239 (1991).

[40] N. N. Bogoljubov, V. V. Tolmachev, and D. V. Shirkov, Fortshr. Phys. **6**, 605 (1958).

[41] J. G. Valatin, Nuovo Cimento **7**, 843 (1958).

[42] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. **108**, 1175 (1957).

[43] H. Wadati, I. Elfimov, and G. A. Sawatzky, Phys. Rev. Lett. **105**, 157004 (2010).

[44] G. Levy, R. Sutarto, D. Chevrier, T. Regier, R. Blyth, J. Geck, S. Wurmehl, L. Harnagea, H. Wadati, T. Mizokawa, I. S. Elfimov, A. Damascelli, and G. A. Sawatzky, Phys. Rev. Lett. **109**, 077001 (2012).